

Letter by Houston et al Regarding Article, “Localized Optogenetic Targeting of Rotors in Atrial Cardiomyocyte Monolayers”

To the Editor:

We read with great interest the article by Feola et al¹ on investigating the mechanisms of rotor-guided ablation through optogenetic stimulation in monolayers of neonatal rat atrial cardiomyocytes. The authors optically map single stable rotors which could be reliably initiated in their homogeneous monolayers and report on the success rate of different light ablation approaches to terminate the rotors by localized conduction block.

We previously studied similar rotor activity using high spatiotemporal resolution optical mapping in monolayers of functionally homogeneous atrial murine cells (HL1-6 myocytes²). At similar mapping resolutions to this work, we also observe apparent spiral waves characteristic of rotor theory. On mapping the cores of our rotors at higher single-cell resolution (2.6×2.6 μm per pixel), we found that the activity was instead driven by microreentry around lines of conduction block/slowing with lengths 200 to 800 μm (<http://www.imperial.ac.uk/imedia/content/view/6049>; Houston C, MRes and Dupont E, PhD, unpublished data, 2017). These lengths correlate with the regularity of the macro spiral wave and contrast the view that the cores are round areas of excitable but unexcited cells as rotor theory postulates.³ Although a different functionally homogeneous biological model, we expect that similar microreentrant mechanisms drive the activity in this study.

The authors reported that their light-based S1-S2 protocol was reliable in initiating stable rotor activity at predetermined locations. Have the authors used their technique to study the core of rotors at higher resolution? In silico studies of spiral waves in neonatal rat atrial cardiomyocyte monolayers suggest that microscopic heterogeneity is necessary for stable rotors.⁴ Our results similarly suggest that microscopic heterogeneity in electrical coupling or cell volume may lead to particular sites being susceptible to microreentry. If this is the case, we would expect rotors to reappear in the same location and with similar dynamics after termination. Was an attempt made to reinitiate a rotor using the same protocol after previous rotor termination? Do new rotors present the same spatial and temporal characteristics observed before termination?

Last, we would like to commend the authors for using a 2-dimensional model of reduced complexity in studying the mechanisms of rotor-guided ablation; we fully support the application of biological models in laying foundations for future research across the translational spectrum.

DISCLOSURES

None.

AFFILIATION

From the ElectroCardioMaths Programme, National Heart and Lung Institute, Imperial College London, United Kingdom.

Charles Houston, MRes
Fu Siong Ng, MRCP, PhD
Emmanuel Dupont, PhD

Circ Arrhythm Electrophysiol is available at <http://circep.ahajournals.org>.

© 2018 American Heart Association, Inc.

REFERENCES

1. Feola I, Volkens L, Majumder R, Teplenin A, Schaliq MJ, Panfilov AV, de Vries AAF, Pijnappels DA. Localized optogenetic targeting of rotors in atrial cardiomyocyte monolayers. *Circ Arrhythm Electrophysiol*. 2017;10:e005591. doi: 10.1161/CIRCEP.117.005591.
2. Dias P, Desplantez T, El-Harasis MA, Chowdhury RA, Ullrich ND, Cabestre-ro de Diego A, Peters NS, Severs NJ, MacLeod KT, Dupont E. Characterisation of connexin expression and electrophysiological properties in stable clones of the HL-1 myocyte cell line. *PLoS One*. 2014;9:e90266. doi: 10.1371/journal.pone.0090266.
3. Pandit SV, Jalife J. Rotors and the dynamics of cardiac fibrillation. *Circ Res*. 2013;112:849–862. doi: 10.1161/CIRCRESAHA.111.300158.
4. Majumder R, Jangsangthong W, Feola I, Ypey DL, Pijnappels DA, Panfilov AV. A mathematical model of neonatal rat atrial monolayers with constitutively active acetylcholine-mediated K⁺ current. *PLoS Comput Biol*. 2016;12:e1004946. doi: 10.1371/journal.pcbi.1004946.

Letter by Houston et al Regarding Article, "Localized Optogenetic Targeting of Rotors in Atrial Cardiomyocyte Monolayers"

Charles Houston, Fu Siong Ng and Emmanuel Dupont

Circ Arrhythm Electrophysiol. 2018;11:

doi: 10.1161/CIRCEP.117.006118

Circulation: Arrhythmia and Electrophysiology is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2018 American Heart Association, Inc. All rights reserved.

Print ISSN: 1941-3149. Online ISSN: 1941-3084

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circep.ahajournals.org/content/11/2/e006118>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation: Arrhythmia and Electrophysiology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation: Arrhythmia and Electrophysiology* is online at:
<http://circep.ahajournals.org/subscriptions/>